

# On block partial linearizations of the pfaffian.

A.A. Lopatin  
 Institute of Mathematics,  
 Siberian Branch of  
 the Russian Academy of Sciences,  
 Pevtsova street, 13,  
 Omsk 644099 Russia  
 artem\_lopatin@yahoo.com  
<http://www.iitam.omsk.net.ru/~lopatin/>

## Abstract

Amitsur's formula, which expresses  $\det(A + B)$  as a polynomial in coefficients of the characteristic polynomial of a matrix, is generalized for partial linearizations of the pfaffian of block matrices. As applications, in upcoming papers we determine generators for the  $SO(n)$ -invariants of several matrices and relations for the  $O(n)$ -invariants of several matrices over a field of arbitrary characteristic.

2000 Mathematics Subject Classification: 15A15.

Key words: pfaffian, partial linearizations.

## 1 Introduction

Throughout,  $K$  is an infinite field of arbitrary characteristic. All vector spaces, algebras, and modules are over  $K$  unless otherwise stated.

A block partial linearization of the pfaffian (b.p.l.p.) is a partial linearization of the pfaffian of block matrices. The concept of b.p.l.p. is important for the invariant theory. Examples of b.p.l.p. include the function DP, of three matrices, used to describe generators for semi-invariants of mixed representations of quivers (see [3]),

the matrix function  $\sigma_{t,s}$ , which describes relations between generators for the invariants of mixed representations of quivers (see [9]), and a partial linearization of the determinant of block matrices as well as the determinant itself (see part 2 of Example 2 for details).

Denote coefficients in the characteristic polynomial of an  $n \times n$  matrix  $X$  by  $\sigma_k(X)$ , i.e.,

$$\det(\lambda E - X) = \lambda^n - \sigma_1(X)\lambda^{n-1} + \cdots + (-1)^n \sigma_n(X).$$

Recall that Amitsur's formula from [1] expresses  $\sigma_k(A+B)$  as a polynomial in  $\sigma_i(a)$ , where  $1 \leq i \leq k$  and  $a$  is a monomial in  $A$  and  $B$  (see Corollary 11 for an explicit formulation). The main result of this paper is the decomposition formula that generalizes Amitsur's formula for a b.p.l.p. and that links a b.p.l.p. with coefficients in the characteristic polynomial of a matrix. This answers the question formulated at the end of Section 4 in [10] as a key consideration for the following problem.

Let  $K_{n,d} = K[x_{ij}(r) \mid 1 \leq i, j \leq n, 1 \leq r \leq d]$  be a polynomial  $K$ -algebra and let  $X_r = (x_{ij}(r))_{1 \leq i, j \leq n}$  be an  $n \times n$  matrix. Note that the subalgebra generated by  $\sigma_k(X_{r_1} \cdots X_{r_s})$  for  $1 \leq k \leq n$  and  $1 \leq r_1, \dots, r_s \leq d$  is the algebra of invariants of several  $n \times n$  matrices (see [2]), and the subalgebra generated by  $\sigma_k(Z_{r_1} \cdots Z_{r_s})$ , where  $Z_r$  is  $X_r$  or its transpose  $X_r^t$  and  $r_1, \dots, r_s$  are in the same range, is the algebra of  $O(n)$ -invariants of several matrices (see [8]). Amitsur's formula plays a crucial role for the description of relations between  $\sigma_k(X_{r_1} \cdots X_{r_s})$  in [7]. The decomposition formula plays the same role for the elements  $\sigma_k(Z_{r_1} \cdots Z_{r_s})$  as it was pointed out in [10]. As a consequence of the decomposition formula we will describe relations for the  $O(n)$ -invariants when the characteristic of  $K$  is different from 2 in the upcoming paper [5].

The decomposition formula will be also used to obtain generators for the  $SO(n)$ -invariants of several matrices when the characteristic of  $K$  is different from 2 (see [4]). Hence we will complete description of generators for the invariants of several matrices, that was originated by Procesi in 1976 (see [6]). Moreover, generators for the invariants under the action of a product of classical groups on the space of (mixed) representations of a quiver will be also established in [4].

An immediate consequence of the decomposition formula is Corollary 12 which states that the product of two pfaffians is a polynomial in  $\sigma_k$ .

The paper is organized as follows.

Section 2 contains general notation.

Section 3 starts with basic definitions of the pfaffian and a b.p.l.p. The key notion of a tableau with substitution is introduced in order to provide a more convenient

language for dealing with b.p.l.p.-s (see Lemma 1). All results are formulated using the notion of a tableau with substitution. Section 3 terminates with the short version of the decomposition formula (see Theorem 2).

The decomposition formula (Theorem 3) and all necessary definitions are stated in Section 4.

The proof of the formula spans the next three sections.

Some applications of the decomposition formula are presented in Section 8.

## 2 Preliminaries

In what follows, all matrices have entries in some commutative unitary algebra over the field  $K$ . Denote the set of non-negative integers by  $\mathbb{N}$ , the set of integers by  $\mathbb{Z}$ , and by  $\mathbb{Q}$  the quotient field of the ring  $\mathbb{Z}$ .

Introduce the lexicographical order  $<$  on  $\mathbb{N} \times \mathbb{N}$  by  $(a_1, b_1) < (a_2, b_2)$  if and only if  $a_1 < a_2$  or  $a_1 = a_2$  and  $b_1 < b_2$ . Denote the composition of functions  $f$  and  $g$  by  $f \circ g(x) = f(g(x))$  or just by  $fg(x)$ .

The cardinality of a set  $S$  is denoted by  $\#S$  and  $S|_{a \rightarrow b}$  denotes the result of substitution of  $b$  for  $a$  in  $S$ . A finite sequence  $a_1, \dots, a_r$  is denoted by  $(a_1, \dots, a_r)$  or just by  $a_1 \cdots a_r$ . In the latter case,  $a_1 \cdots a_r$  is called a word in letters  $a_1, \dots, a_r$ . Denote the degree of a word  $w$  in a letter  $a$  by  $\deg_a w$ . Similarly to the cardinality of a set, we denote the length of a sequence  $\underline{\lambda} = (\lambda_1, \dots, \lambda_p)$  by  $\#\underline{\lambda} = p$ . We use notation  $\{\dots\}_m$  for multisets, i.e., given an equivalence  $=$  on a set  $S$  and  $a_1, \dots, a_p, b_1, \dots, b_q \in S$ , we write  $\{a_1, \dots, a_p\}_m = \{b_1, \dots, b_q\}_m$  if and only if  $p = q$  and

$$\#\{1 \leq j \leq p \mid a_j = a_i\} = \#\{1 \leq j \leq p \mid b_j = a_i\}$$

for any  $1 \leq i \leq p$ . For integers  $i < j$  denote the interval  $i, i+1, \dots, j-1, j$  by  $[i, j]$ .

Denote by  $id_n$  the identity permutation of the group  $\mathcal{S}_n$  of permutations on  $n$  elements. Given  $i \neq j$ ,  $(i, j)$  is the transposition interchanging  $i$  and  $j$ .

An element  $u$  of a free monoid is called *primitive* if  $u \neq u_0 \cdots u_0$ , where the number of factors is more than 1.

A vector  $\underline{\lambda} = (\lambda_1, \dots, \lambda_l) \in \mathbb{N}^l$  satisfying  $\lambda_1 \geq \dots \geq \lambda_l$  and  $\lambda_1 + \dots + \lambda_l = t$  is called a *partition* of  $t$  and is denoted by  $\underline{\lambda} \vdash t$ . A *multi-partition*  $\underline{\lambda} \vdash \underline{t}$  is a  $q$ -tuple of partitions  $\underline{\lambda} = (\underline{\lambda}_1, \dots, \underline{\lambda}_q)$ , where  $\underline{\lambda}_i \vdash t_i$ ,  $\underline{t} = (t_1, \dots, t_q) \in \mathbb{N}^q$ .

## 3 Definitions

Let us recall definitions of the pfaffian and its partial linearizations.

If  $n$  is even, then the pfaffian of an  $n \times n$  skew-symmetric matrix  $X = (x_{ij})$  is

$$\text{pf}(X) = \sum \text{sgn}(\pi) \prod_{i=1}^{n/2} x_{\pi(2i-1), \pi(2i)},$$

where the sum ranges over permutations  $\pi \in \mathcal{S}_n$  satisfying  $\pi(1) < \pi(3) < \dots < \pi(n-1)$  and  $\pi(2i-1) < \pi(2i)$  for all  $1 \leq i \leq n/2$ . Define the *generalized pfaffian* of an arbitrary  $n \times n$  matrix  $X = (x_{ij})$  by

$$\overline{\text{pf}}(X) = \text{pf}(X - X^t).$$

By abuse of notation we will refer to  $\overline{\text{pf}}$  as the pfaffian. For  $K = \mathbb{Q}$  there is a more convenient formula

$$\overline{\text{pf}}(X) = \text{pf}(X - X^t) = \frac{1}{(n/2)!} \sum_{\pi \in \mathcal{S}_n} \text{sgn}(\pi) \prod_{i=1}^{n/2} x_{\pi(2i-1), \pi(2i)}. \quad (1)$$

Note that for an  $n \times n$  skew-symmetric matrix  $X = (x_{ij})$  we have

$$\overline{\text{pf}}(X) = 2^{n/2} \text{pf}(X).$$

For  $n \times n$  matrices  $X_1 = (x_{ij}(1)), \dots, X_s = (x_{ij}(s))$  and positive integers  $k_1, \dots, k_s$ , satisfying  $k_1 + \dots + k_s = n/2$ , consider the polynomial  $\overline{\text{pf}}(x_1 X_1 + \dots + x_s X_s)$  in the variables  $x_1, \dots, x_s$ . The partial linearization  $\overline{\text{pf}}_{k_1, \dots, k_s}(X_1, \dots, X_s)$  of the pfaffian is the coefficient of  $x_1^{k_1} \dots x_s^{k_s}$  in this polynomial. In other words, for  $K = \mathbb{Q}$  we have

$$\overline{\text{pf}}_{k_1, \dots, k_s}(X_1, \dots, X_s) = \frac{1}{c} \sum_{\pi \in \mathcal{S}_n} \text{sgn}(\pi) \prod_{j=1}^s \prod_{i=k_1 + \dots + k_{j-1} + 1}^{k_1 + \dots + k_j} x_{\pi(2i-1), \pi(2i)}(j), \quad (2)$$

where  $c = k_1! \dots k_s!$ . The partial linearization  $\det_{k_1, \dots, k_s}(X_1, \dots, X_s)$  of the determinant is defined analogously, where  $X_1, \dots, X_s$  are  $n \times n$  matrices,  $k_1 + \dots + k_s = n$ , and  $n$  is arbitrary.

Now we can give the definition of a b.p.l.p.

Fix  $\underline{n} = (n_1, \dots, n_m) \in \mathbb{N}^m$ , where  $n = n_1 + \dots + n_m$  is even. For any  $1 \leq p, q \leq m$  and an  $n_p \times n_q$  matrix  $X$  denote by  $X^{p,q}$  the  $n \times n$  matrix, partitioned into  $m \times m$  number of blocks, where the block in the  $(i, j)$ -th position is an  $n_i \times n_j$  matrix; the block in the  $(p, q)$ -th position is equal to  $X$ , and the rest of blocks are zero matrices.

Let  $1 \leq p_1, \dots, p_s, q_1, \dots, q_s \leq m$ , let  $X_j$  be an  $n_{p_j} \times n_{q_j}$  matrix for any  $1 \leq j \leq s$ , and let  $k_1, \dots, k_s$  be positive integers, satisfying  $k_1 + \dots + k_s = n/2$ . The element

$$\overline{\text{pf}}_{k_1, \dots, k_s}(X_1^{p_1, q_1}, \dots, X_s^{p_s, q_s}) \quad (3)$$

is a partial linearization of the pfaffian of block matrices  $X_1^{p_1, q_1}, \dots, X_s^{p_s, q_s}$ , and it is called a *block partial linearization of the pfaffian (b.p.l.p.)*. We consider only b.p.l.p.-s, satisfying

$$\sum_{1 \leq j \leq s, p_j=i} k_j + \sum_{1 \leq j \leq s, q_j=i} k_j = n_i \text{ for all } 1 \leq i \leq m, \quad (4)$$

since only these elements appear in the invariant theory in the context of Section 1.

**Example 1.** Let  $\underline{n} = (5, 3)$  and let  $X_1, X_2, X_3$  be  $5 \times 3$ ,  $5 \times 5$ , and  $3 \times 3$  matrices respectively. Then

$$X_1^{1,2} = \begin{pmatrix} 0 & X_1 \\ 0 & 0 \end{pmatrix}, \quad X_2^{1,1} = \begin{pmatrix} X_2 & 0 \\ 0 & 0 \end{pmatrix}, \quad X_3^{2,2} = \begin{pmatrix} 0 & 0 \\ 0 & X_3 \end{pmatrix}$$

are  $8 \times 8$  matrices. The b.p.l.p.  $f = \overline{\text{pf}}_{1,2,1}(X_1^{1,2}, X_2^{1,1}, X_3^{2,2})$  satisfies condition (4). It is convenient to display the data that determine  $f$  as a two-column tableau filled with arrows:

|              |               |
|--------------|---------------|
| $a$          | $\rightarrow$ |
| $b$          | $d$           |
| $\downarrow$ | $\downarrow$  |
| $c$          |               |
| $\downarrow$ |               |

The arrow  $a$  determines the block matrix  $X_1^{1,2}$  as follows:  $a$  goes from the 1st column to the 2nd column and we assign the matrix  $X_1$  to  $a$ . To determine the block matrix  $X_2^{1,1}$  we take *two* arrows  $b, c$  that go from the 1st column to the 1st column, since the degree of  $f$  in entries of  $X_2$  is 2. Assign the matrix  $X_2$  to  $b$  and to  $c$ . Finally, the arrow  $d$  determines the block matrix  $X_3^{2,2}$ :  $d$  goes from the 2nd column to the 2nd column and  $X_3$  is assigned to it. Condition (4) implies that for every column the total amount of arrows that start or terminate in the column is equal to the length of the column. Note that this conditions do not uniquely determine arrows of a tableau.

Below we formulate the definition of a tableau with substitution that gives alternative way to work with b.p.l.p.-s. Acting in the same way as in Example 1, for every


b.p.l.p.  $f$  given by (3) we construct a tableau with substitution  $(T, (X_1, \dots, X_s))$  and define the function  $\text{bpf}_T$  such that  $f = \pm \text{bpf}_T(X_1, \dots, X_s)$ .

**Definition (of shapes).** A *shape* of dimension  $\underline{n} = (n_1, \dots, n_m) \in \mathbb{N}^m$  is a collection of  $m$  columns of cells. The columns are numbered by  $1, 2, \dots, m$ , and the  $i$ -th column contains exactly  $n_i$  cells, where  $1 \leq i \leq m$ . Numbers  $1, \dots, n_i$  are assigned to the cells of the  $i$ -th column, starting from the top. As an example, the shape of dimension  $\underline{n} = (3, 2, 3, 1, 1)$  is

| 1 | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| 1 | 1 | 1 | 1 | 1 |
| 2 | 2 | 2 |   |   |
| 3 |   | 3 |   |   |

**Definition (of a tableau with substitution).** Let  $\underline{n} = (n_1, \dots, n_m) \in \mathbb{N}^m$  and let  $n = n_1 + \dots + n_m$  be even. A pair  $(T, (X_1, \dots, X_s))$  is called a *tableau with substitution* of dimension  $\underline{n}$  if

- $T$  is a shape of dimension  $\underline{n}$  that is filled with arrows. An *arrow* goes from one cell of the shape into another one, and each cell of the shape is either the head or the tail of one and only one arrow. We refer to  $T$  as a *tableau* of dimension  $\underline{n}$ , and we write  $a \in T$  for an arrow  $a$  from  $T$ . Given an arrow  $a \in T$ , denote by  $a'$  and  $a''$  the columns containing the head and the tail of  $a$ , respectively. Similarly, denote by  $'a$  the number assigned to the cell containing the head of  $a$ , and denote by  $''a$  the number assigned to the cell containing the tail of  $a$ . Schematically this is depicted as



- $(X_1, \dots, X_s)$  is a sequence of matrices, where the  $(p, q)$ -th entry of  $X_j$  is  $(X_j)_{pq}$  ( $1 \leq j \leq s$ ). We assume that an  $n_{a''} \times n_{a'}$  matrix  $X_{\varphi(a)}$  is assigned to an arrow  $a \in T$ , where  $1 \leq \varphi(a) \leq s$ , and  $\{\varphi(a) \mid a \in T\} = [1, s]$ . Moreover, we assume that if  $a, b \in T$  and  $\varphi(a) = \varphi(b)$ , then  $a' = b'$ ,  $a'' = b''$ .

**Definition (of  $\text{bpf}_T(X_1, \dots, X_s)$ ).** Let  $(T, (X_1, \dots, X_s))$  be a tableau with substitution of dimension  $\underline{n}$ . Define

$$\text{bpf}_T(X_1, \dots, X_s) = \sum \text{sgn}(\pi_1) \cdots \text{sgn}(\pi_m) \prod_{a \in T} (X_{\varphi(a)})_{\pi_{a''}('a), \pi_{a'}('a)},$$

where the sum ranges over permutations  $\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_m \in \mathcal{S}_{n_m}$  such that for any  $a, b \in T$  the conditions  $\varphi(a) = \varphi(b)$  and  $''a < ''b$  imply that  $\pi_i(''a) < \pi_i(''b)$  for  $i = a'' = b''$ . Note that each coefficient in  $\text{bpf}_T(X_1, \dots, X_s)$ , considered as a polynomial in entries of  $X_1, \dots, X_s$ , is  $\pm 1$ . For  $K = \mathbb{Q}$  there is a more convenient formula

$$\text{bpf}_T(X_1, \dots, X_s) = \frac{1}{c_T} \text{bpf}_T^0(X_1, \dots, X_s),$$

where

$$\text{bpf}_T^0(X_1, \dots, X_s) = \sum_{\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_m \in \mathcal{S}_{n_m}} \text{sgn}(\pi_1) \cdots \text{sgn}(\pi_m) \prod_{a \in T} (X_{\varphi(a)})_{\pi_{a''}(''a), \pi_{a'}(''a)}, \quad (5)$$

and the coefficient  $c_T$  is equal to

$$\prod_{j=1}^s \#\{a \in T \mid \varphi(a) = j\}!.$$

**Lemma 1.**

- a) Let  $f$  be a b.p.l.p. (3) satisfying (4). Then there is a tableau with substitution  $(T, (X_1, \dots, X_s))$  such that  $\text{bpf}_T(X_1, \dots, X_s) = \pm f$ .
- b) For every tableau with substitution  $(T, (X_1, \dots, X_s))$  of dimension  $\underline{n}$  there is a b.p.l.p.  $f$  satisfying (4) such that  $\text{bpf}_T(X_1, \dots, X_s) = \pm f$ .


*Proof.* **a)** A tableau with substitution  $(T, (X_1, \dots, X_s))$  of dimension  $\underline{n}$  is constructed as follows. Its arrows are  $a_{jr}$ , where  $a_{jr}$  goes from the  $p_j$ -th to the  $q_j$ -th column for  $1 \leq j \leq s$  and  $1 \leq r \leq k_j$ . Condition (4) guarantees that for any  $1 \leq i \leq m$  the total number of arrows that begin or end in the  $i$ -th column is  $n_i$ . Complete the construction by setting  $\varphi(a_{jr}) = j$ . Note that  $(T, (X_1, \dots, X_s))$  is not uniquely determined by  $f$ .

**b)** Consider  $a_1, \dots, a_s \in T$  such that  $\varphi(a_1) = 1, \dots, \varphi(a_s) = s$ . Then

$$\text{bpf}_T(X_1, \dots, X_s) = \pm \overline{\text{pf}}_{k_1, \dots, k_s}(X_1^{a_1'', a_1'}, \dots, X_s^{a_s'', a_s'}),$$

where  $k_j = \#\{a \in T \mid \varphi(a) = j\}$  for any  $1 \leq j \leq s$ .  $\square$

**Example 2. 1.** Let  $n$  be even and  $(T, (X_1, \dots, X_s))$  be a tableau with substitution of dimension  $(n)$ , where  $T$  is



In other words, arrows of  $T$  are  $a_1, \dots, a_{n/2}$ , where  $'a_i = 2i$ ,  $''a_i = 2i - 1$ ,  $a'_i = a''_i = 1$  for  $1 \leq i \leq n/2$ , and  $X_1, \dots, X_s$  are  $n \times n$  matrices.


If  $s = 1$ , then (1) implies  $\text{bpf}_T(X_1) = \overline{\text{pf}}(X_1)$  for  $K = \mathbb{Q}$  and consequently for an arbitrary  $K$ .

If  $s > 1$ , then by (2)

$$\text{bpf}_T(X_1, \dots, X_s) = \overline{\text{pf}}_{k_1, \dots, k_s}(X_1, \dots, X_s),$$

where  $k_j = \#\{a \in T \mid \varphi(a) = j\}$  for any  $1 \leq j \leq s$ , is a partial linearization of the pfaffian.

**2.** For  $n \times n$  matrices  $X_1, \dots, X_s$  let  $(T, (X_1, \dots, X_s))$  be the tableau with substitution of dimension  $(n, n)$ , where  $T$  is



If  $s = 1$ , then the formula

$$\det(X) = \frac{1}{n!} \sum_{\pi_1, \pi_2 \in \mathcal{S}_n} \text{sgn}(\pi_1) \text{sgn}(\pi_2) \prod_{i=1}^n x_{\pi_1(i), \pi_2(i)}, \quad (6)$$

which is valid over  $\mathbb{Q}$ , implies the equality  $\text{bpf}_T(X_1) = \det(X_1)$  over every  $K$ . For  $s > 1$  the expression  $\text{bpf}_T(X_1, \dots, X_s)$  is a partial linearization of the determinant.

**3.** If  $(T, (X_1, X_2))$  is a tableau with substitution from part 2, where for  $1 \leq k \leq n$  we have  $\varphi(a_1) = \dots = \varphi(a_k) = 1$ ,  $\varphi(a_{k+1}) = \dots = \varphi(a_n) = 2$ ,  $X_1 = X$ , and  $X_2 = E$  is the identity  $n \times n$  matrix, then  $\text{bpf}_T(X, E) = \sigma_k(X)$ .

**4.** Suppose that  $t, s, r \in \mathbb{N}$ , and  $X, Y, Z$ , respectively, are matrices of dimensions  $(t + 2s) \times (t + 2r)$ ,  $(t + 2s) \times (t + 2s)$ ,  $(t + 2r) \times (t + 2r)$ , respectively. Let  $T$  be a

tableau of dimension  $(t + 2s, t + 2r)$  that in the case  $s = r$  is depicted as

|              |               |
|--------------|---------------|
| $a_1$        | $\rightarrow$ |
| $\vdots$     | $\vdots$      |
| $a_t$        | $\rightarrow$ |
| $b_1$        | $c_1$         |
| $\downarrow$ | $\downarrow$  |
| $\vdots$     | $\vdots$      |
| $b_s$        | $c_r$         |
| $\downarrow$ | $\downarrow$  |

otherwise define  $T$  analogously. Let  $(T, (X, Y, Z))$  be a tableau with substitution, where  $\varphi(a_i) = 1$ ,  $\varphi(b_j) = 2$ , and  $\varphi(c_k) = 3$  for  $1 \leq i \leq t$ ,  $1 \leq j \leq s$ , and  $1 \leq k \leq r$ . Then  $\text{bpf}_T(X, Y, Z) = \pm \text{DP}_{s,r}(X, Y, Z)$ , where  $\text{DP}_{s,r}$  was introduced in Section 3 of [3].

The main result of the paper is the following decomposition formula.

**Theorem 2. (Decomposition formula: short version)**

Let  $(T, (X_1, \dots, X_s))$  be a tableau with substitution of dimension  $\underline{n} \in \mathbb{N}^m$ . Let  $1 \leq q_1 < q_2 \leq m$ ,  $n_{q_1} = n_{q_2}$ , and the vector  $\underline{d} \in \mathbb{N}^{m-2}$  be obtained from  $\underline{n}$  by eliminating the  $q_1$ -th and the  $q_2$ -th coordinates. Then  $\text{bpf}_T(X_1, \dots, X_s)$  is a polynomial in  $\text{bpf}_D(Y_1, \dots, Y_l)$  and  $\sigma_k(h)$ , where

- $(D, (Y_1, \dots, Y_l))$  ranges over tableaux with substitutions of dimension  $\underline{d}$  such that  $Y_1, \dots, Y_l$  are products of matrices  $X_1, \dots, X_s, X_1^t, \dots, X_s^t$ ;
- $h$  ranges over products of matrices  $X_1, \dots, X_s, X_1^t, \dots, X_s^t$ ;
- $k$  ranges over  $[1, n_{q_1}]$ .

Moreover, coefficients of this polynomial belong to the image of  $\mathbb{Z}$  in  $K$  under the natural homomorphism.

## 4 Decomposition formula

Consider a tableau with substitution  $(T, (X_1, \dots, X_s))$  of dimension  $\underline{n} \in \mathbb{N}^m$ . In what follows we assume that  $n_{q_1} = n_{q_2} = n$  for  $1 \leq q_1 < q_2 \leq m$ .

Denote by  $\mathcal{M}$  the monoid freely generated by letters  $x_1, x_2, \dots, x_1^t, x_2^t, \dots$ . Fix some lexicographical order  $<$  on  $\mathcal{M}$ . Let  $\mathcal{M}_T$  be the submonoid of  $\mathcal{M}$ , generated by  $x_1, \dots, x_s, x_1^t, \dots, x_s^t$ . For short, we will write  $1, \dots, s, 1^t, \dots, s^t$  instead of  $x_1, \dots, x_s, x_1^t, \dots, x_s^t$ . Given  $a \in T$ , we consider  $\varphi(a) \in \{1, \dots, s\}$  as an element of  $\mathcal{M}_T$ .

For  $u \in \mathcal{M}_T$  define the matrix  $X_u$  by the following rules:

- $X_{j^t} = X_j^t$  for any  $1 \leq j \leq s$ ;
- $X_{vw} = \begin{cases} X_v X_w, & \text{if the product of these matrices is well defined} \\ 0, & \text{otherwise} \end{cases}$   
for  $v, w \in \mathcal{M}_T$ .

For an arrow  $a \in T$  denote by  $a^t$  the *transpose arrow*, i.e., by definition  $(a^t)'' = a'$ ,  $(a^t)' = a''$ ,  $''(a^t) = 'a$ ,  $'(a^t) = ''a$ ,  $\varphi(a^t) = \varphi(a)^t \in \mathcal{M}_T$ . Obviously,  $(a^t)^t = a$ .

We write  $a \stackrel{t}{\in} T$  if  $a \in T$  or  $a^t \in T$ .

**Definition (of paths).** We say that  $a_1, a_2 \stackrel{t}{\in} T$  are *successive* in  $T$ , if  $a'_1, a''_2 \in \{q_1, q_2\}$ ,  $a'_1 \neq a''_2$ ,  $'a_1 = ''a_2$ .

A word  $a = a_1 \cdots a_r$ , where  $a_1, \dots, a_r \stackrel{t}{\in} T$ , is called a *path* in  $T$  with respect to columns  $q_1$  and  $q_2$ , if  $a_i, a_{i+1}$  are successive for any  $1 \leq i \leq r-1$ . In this case by definition  $\varphi(a) = \varphi(a_1) \cdots \varphi(a_r) \in \mathcal{M}_T$  and  $a^t = a_r^t \cdots a_1^t$  is a path in  $T$ ; we denote  $a'_r, 'a_r, a''_1, ''a_1$ , respectively, by  $a', 'a, a'', ''a$ , respectively. Since the columns  $q_1$  and  $q_2$  are fixed, we usually refer to  $a$  as a path in  $T$ .

A path  $a_1 \cdots a_r$  is *open* if both  $a''_1, a'_r \notin \{q_1, q_2\}$ .

A path  $a_1 \cdots a_r$  is *closed* if  $a_r, a_1$  are successive; in particular,  $a''_1, a'_r \in \{q_1, q_2\}$ .

An element  $a \stackrel{t}{\in} T$  with  $a', a'' \in \{q_1, q_2\}$  and  $'a = ''a$  is also called a closed path.

A path is called *maximal* if it is either open or closed.

**Example 3.** Let



be fragments of two tableaux, where in both cases we have drawn columns  $q_1, q_2$ . In the first case we have an open path  $a^t b^t c d$ ; and in the second case we have closed paths:  $a, b c^t$ .

Consider words  $a = a_1 \cdots a_p$ ,  $b = b_1 \cdots b_q$ , where  $a_1 \cdots a_p, b_1 \cdots b_q \in T$  (in particular, both  $a, b$  might be paths in  $T$ ). We write

- $a = b$  if  $p = q$  and  $a_1 = b_1, \dots, a_p = b_p$ ;
- $a \stackrel{t}{=} b$  if  $a = b$  or  $a = b^t$ ;
- $a \stackrel{c}{=} b$  if there is a cyclic permutation  $\pi \in \mathcal{S}_p$  such that  $a_{\pi(1)} \cdots a_{\pi(p)} = b$ ;
- $a \stackrel{ct}{=} b$  if there is a cyclic permutation  $\pi \in \mathcal{S}_p$  such that  $a_{\pi(1)} \cdots a_{\pi(p)} \stackrel{t}{=} b$ .

Note that  $\stackrel{t}{=}$ ,  $\stackrel{c}{=}$ , and  $\stackrel{ct}{=}$  are equivalences. We will use similar notations for elements of  $\mathcal{M}_T$ , for sets of paths, and for subsets of  $\mathcal{M}_T$ . In the same fashion define the signs of inclusion  $\stackrel{t}{\in}$ ,  $\stackrel{ct}{\in}$ , and the sign of subtraction of sets  $\stackrel{t}{\setminus}$ . As an example, if  $s \geq 6$ , then we have the following equivalence of subsets of  $\mathcal{M}_T$

$$\{12^t, 3^t4, 56\} \stackrel{t}{\setminus} \{21^t\} \stackrel{t}{=} \{4^t3, 56\}.$$

Note that

- if  $a$  is an open path and  $a \stackrel{t}{=} b$ , then  $b$  is also an open path;
- if  $a$  is a closed path and  $a \stackrel{ct}{=} b$ , then  $b$  is also a closed path;
- if  $a = a_1 \cdots a_r$ , where  $a_1, \dots, a_r \stackrel{t}{\in} T$ , is an open path, then  $\varphi(a) \in \mathcal{M}_T$  is a primitive element (see Section 2 for the definition), since otherwise there is an  $1 < l \leq r$  such that  $\varphi(a_1) = \varphi(a_l)$ ; hence  $a_1'' = a_l''$ ; on the other hand,  $a_1'' \notin \{q_1, q_2\}$  and  $a_l'' \in \{q_1, q_2\}$ ; a contradiction.

Denote by  $T_o$  the set  $\{b \mid b \text{ is an open path in } T, (b'', ''b) < (b', 'b)\}$ , where  $<$  is the lexicographical order on  $\mathbb{N} \times \mathbb{N}$  introduced in Section 2. Obviously,  $T_o$  is a set of representatives of open paths in  $T$  with respect to the  $\stackrel{t}{=}$ -equivalence, i.e.,  $a \stackrel{t}{\in} T_o$  for every open path  $a$  in  $T$ , and  $a \stackrel{t}{\neq} b$  for all  $a, b \in T_o$  with  $a \neq b$ . Let  $T_{cl}$  be a set of representatives of closed paths in  $T$  with respect to the  $\stackrel{ct}{=}$ -equivalence. Define

$$\varphi(T_o) = \{\varphi(a) \mid a \in T_o\}_m, \quad \varphi(T_{cl}) = \{\varphi(a) \mid a \in T_{cl}\}_m$$

(see Section 2 for the definition of a multiset). Then the expression

$$\text{tr}(T_{cl}) = \prod_{a \in T_{cl}} \text{tr}(X_{\varphi(a)}).$$

is well defined.

**Definition (of  $T^\tau$  and  $\tilde{T}$ ).** **a)** Let  $\tau \in \mathcal{S}_n$ . Permute the cells of the  $q_2$ -th column of  $T$  by  $\tau$  and denote the resulting tableau by  $T^\tau$ . The arrows of  $T^\tau$  are  $\{a^\tau \mid a \in T\}$ , where  $\varphi(a^\tau) = \varphi(a)$ ,  $(a^\tau)'' = a''$ ,  $(a^\tau)' = a'$ , and

$$''(a^\tau) = \begin{cases} \tau('a), & \text{if } a'' = q_2 \\ ''a, & \text{otherwise} \end{cases}, \quad '(a^\tau) = \begin{cases} \tau('a), & \text{if } a' = q_2 \\ 'a, & \text{otherwise} \end{cases}.$$

Obviously,  $(T^\tau, (X_1, \dots, X_s))$  is a tableau with substitution.

**b)** Define the tableau with substitution  $(\tilde{T}, X_{\tilde{T}})$  of dimension  $(n_1, \dots, n_{q_1-1}, n_{q_1+1}, \dots, n_{q_2-1}, n_{q_2+1}, \dots, n_m)$ , where  $X_{\tilde{T}} = (Y_1, \dots, Y_t)$  for some matrices  $Y_1, \dots, Y_t$  for suitable  $t > 0$  as follows. If  $q_1 = m - 1$ ,  $q_2 = m$ , then  $\tilde{T}$  contains two types of arrows:

- $\tilde{a}$ , where  $a \in T$ , and both  $a', a'' \notin \{q_1, q_2\}$ ;
- $\tilde{a}$ , where  $a$  is an open path from  $T_o$ .

In both cases the tail and the head of  $\tilde{a}$  coincides with the tail and the head of  $a$ , respectively. Define  $t > 0$  and  $\varphi(u)$  for  $u \in \tilde{T}$  in such a way that  $\{\varphi(u) \mid u \in \tilde{T}\} = [1, t]$  and for all  $\tilde{b}, \tilde{c} \in \tilde{T}$  we have

- the equality  $\varphi(\tilde{b}) = \varphi(\tilde{c})$  holds if and only if  $\varphi(b) = \varphi(c)$ ,
- the inequality  $\varphi(\tilde{b}) < \varphi(\tilde{c})$  holds if and only if  $\varphi(b) < \varphi(c)$ ,

where  $<$  is the lexicographical order on  $\mathcal{M}$  fixed at the beginning of this section. We set  $Y_{\varphi(\tilde{a})} = X_{\varphi(a)}$ . Then the tableau with substitution  $(\tilde{T}, X_{\tilde{T}})$  is well defined.

For arbitrary  $q_1, q_2$  the definition of  $(\tilde{T}, X_{\tilde{T}})$  is obtained analogously using an appropriate shift to define numbers  $\tilde{a}', \tilde{a}''$ .

For short, we will omit parentheses and use the following conventions

$$\begin{aligned} (T^\tau)_o &= T_o^\tau, & (T^\tau)_{cl} &= T_{cl}^\tau, \\ \widetilde{(T^\tau)} &= \tilde{T}^\tau, & \widetilde{(a^\tau)} &= \tilde{a}^\tau, \\ \widetilde{(T^\tau)}_o &= \tilde{T}_o^\tau, & \widetilde{(T^\tau)}_{cl} &= \tilde{T}_{cl}^\tau, \end{aligned}$$

for any  $\tau \in \mathcal{S}_n$  and  $a \in T$ .

**Example 4. 1.** Let  $T$  be the tableau 

|     |               |              |
|-----|---------------|--------------|
| $a$ | $\rightarrow$ | $b$          |
| $c$ | $\rightarrow$ | $\downarrow$ |

 of dimension  $(2, 2, 2)$ . Suppose  $q_1 = 2$  and  $q_2 = 3$ . Then  $\tilde{T}$  is 

|              |
|--------------|
| $d$          |
| $\downarrow$ |

, where  $d = \widetilde{abc^t}$ .

2. Let  $T$  be the tableau

|              |            |               |
|--------------|------------|---------------|
| $a$          | $b$        | $\rightarrow$ |
| $\downarrow$ |            |               |
| $c$          | $\nearrow$ | $\nearrow$    |
|              | $d$        |               |

of dimension  $(3, 3, 2)$ . Suppose  $q_1 = 1$ ,  $q_2 = 2$ , and  $\tau \in \mathcal{S}_3$  is the cycle  $(1, 2, 3)$ . Then  $T^\tau$  is

|              |               |            |
|--------------|---------------|------------|
| $a^\tau$     | $d^\tau$      | $\nearrow$ |
| $\downarrow$ | $b^\tau$      | $\searrow$ |
| $c^\tau$     | $\rightarrow$ |            |

$\tilde{T}$  is  $\begin{bmatrix} e \\ \downarrow \end{bmatrix}$ , and  $\tilde{T}^\tau$  is  $\begin{bmatrix} h \\ \downarrow \end{bmatrix}$ , where  $e = \widetilde{b^t a c^t d}$ ,  $h = \widetilde{(b^\tau)^t (a^\tau)^t d^\tau}$ . In particular,  $\tilde{T}$  and  $\tilde{T}^\tau$  are different.

**Definition (of admissibility).** Let  $\underline{\beta} \in \mathbb{N}^p$ ,  $\underline{\gamma} \in \mathbb{N}^q$ ,  $\underline{b} = (b_1, \dots, b_p)$ ,  $\underline{c} = (c_1, \dots, c_q)$ , where  $b_i, c_j$  are primitive elements of  $\mathcal{M}_T$ . Moreover, let  $b_1, \dots, b_p$  be pairwise different with respect to  $\stackrel{t}{=}$ , i.e.,  $b_i \stackrel{t}{\neq} b_j$  for  $i \neq j$ ; and let  $c_1, \dots, c_q$  be pairwise different with respect to  $\stackrel{ct}{=}$ . Then  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$  is called a  $T$ -quadruple.

A  $T$ -quadruple  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$  is called  $T$ -admissible if for some  $\xi = \xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}} \in \mathcal{S}_n$  the following equivalences of multisets hold:

$$\varphi(T_o^\xi) \stackrel{t}{=} \underbrace{\{b_1, \dots, b_1\}}_{\beta_1}, \dots, \underbrace{\{b_p, \dots, b_p\}}_{\beta_p}, \quad \varphi(T_{cl}^\xi) \stackrel{ct}{=} \underbrace{\{c_1, \dots, c_1\}}_{\gamma_1}, \dots, \underbrace{\{c_q, \dots, c_q\}}_{\gamma_q}.$$

We write  $(\underline{\beta}^0, \underline{\gamma}^0, \underline{b}^0, \underline{c}^0) \stackrel{ct}{=} (\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$  and say that these quadruples are equivalent if and only if

$$\begin{aligned} \underbrace{\{b_1^0, \dots, b_1^0\}}_{\beta_1^0}, \dots, \underbrace{\{b_p^0, \dots, b_p^0\}}_{\beta_p^0} &\stackrel{t}{=} \underbrace{\{b_1, \dots, b_1\}}_{\beta_1}, \dots, \underbrace{\{b_p, \dots, b_p\}}_{\beta_p}, \\ \underbrace{\{c_1^0, \dots, c_1^0\}}_{\gamma_1^0}, \dots, \underbrace{\{c_q^0, \dots, c_q^0\}}_{\gamma_q^0} &\stackrel{ct}{=} \underbrace{\{c_1, \dots, c_1\}}_{\gamma_1}, \dots, \underbrace{\{c_q, \dots, c_q\}}_{\gamma_q}. \end{aligned}$$

If  $(\underline{\beta}^0, \underline{\gamma}^0, \underline{b}^0, \underline{c}^0) \stackrel{ct}{=} (\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$  and  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$  is  $T$ -admissible, then the quadruple  $(\underline{\beta}^0, \underline{\gamma}^0, \underline{b}^0, \underline{c}^0)$  also has the same property, since we can take  $\xi_{\underline{\beta}^0, \underline{\gamma}^0, \underline{b}^0, \underline{c}^0} = \xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ .

Denote by  $Q_T$  a set of representatives of  $T$ -admissible quadruples with respect to the  $\stackrel{ct}{=}$ -equivalence.

**Theorem 3. (Decomposition formula).** *Let  $1 \leq q_1 < q_2 \leq m$  and  $n_{q_1} = n_{q_2} = n$ . Then for a tableau with substitution  $(T, (X_1, \dots, X_s))$  of dimension  $\underline{n}$  we have*

$$\text{bpf}_T(X_1, \dots, X_s) = \sum_{(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T} \text{sgn}(\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}) \text{bpf}_{\tilde{T}^{\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}}}(X_{\tilde{T}^{\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}}}) \prod_{j=1}^{\#\underline{\gamma}} \sigma_{\gamma_j}(X_{c_j}).$$

Neither the permutation  $\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$  nor its sign are unique for a representative of the  $\stackrel{ct}{=}$ -equivalence class of a  $T$ -admissible quadruple  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$ . However, it follows from part b) of Lemma 9, which is formulated below, that  $\text{sgn}(\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}) \text{bpf}_{\tilde{T}^{\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}}}(X_{\tilde{T}^{\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}}})$  does not depend on a representative of the  $\stackrel{ct}{=}$ -equivalence class. Thus the right hand side of the decomposition formula is well defined.

Sections 5, 6, and 7 are devoted to a proof of this theorem. Since the right hand side of the decomposition formula is a polynomial with integer coefficients in  $\mathbb{Z}$ , if the characteristic of  $K$  is zero, or in  $\mathbb{Z}/p\mathbb{Z}$ , if the characteristic of  $K$  is  $p > 0$ ; it is enough to prove the theorem in the case  $K = \mathbb{Q}$ . To simplify notation we assume that  $q_1 = m - 1$ ,  $q_2 = m$ .

The proof is organized as follows. In Section 5 we prove Lemma 4 for  $\text{bpf}_T^0(X_1, \dots, X_s)$ , which is a statement analogous to Theorem 3. The short version of the decomposition formula (see Theorem 2) follows from Lemma 4 in the case of characteristic zero. In order to prove Theorem 2 as well as Theorem 3 over the field of arbitrary characteristic we, working over  $\mathbb{Q}$ , should divide the formula in Lemma 4 by  $c_T$ . This problem is solved by rewriting  $\text{bpf}_T(X_1, \dots, X_s)$  in a more suitable form (see formula (14) in Section 6) and by applying Lemma 10 from Section 7.

## 5 Decomposition of $\text{bpf}_T^0(X_1, \dots, X_s)$

**Lemma 4.** *Let  $\underline{n} \in \mathbb{N}^m$ ,  $1 \leq q_1 < q_2 \leq m$ ,  $n_{q_1} = n_{q_2} = n$ , and  $(T, (X_1, \dots, X_s))$  be a tableau with substitution of dimension  $\underline{n}$ . Then*

$$\text{bpf}_T^0(X_1, \dots, X_s) = \sum_{\tau \in \mathcal{S}_n} \text{sgn}(\tau) \text{bpf}_{\tilde{T}^\tau}^0(X_{\tilde{T}^\tau}) \text{tr}(T_{cl}^\tau).$$

*Proof.* Without loss of generality we can assume that  $q_1 = m - 1$ ,  $q_2 = m$ . Define

$$A = \{a \in T \mid a' > m - 2 \text{ or } a'' > m - 2\}.$$

For fixed  $\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_{m-2} \in \mathcal{S}_{n_{m-2}}$  define

$$f_0 = \sum_{\pi_{m-1}, \pi_m \in \mathcal{S}_n} \text{sgn}(\pi_{m-1}) \text{sgn}(\pi_m) \prod_{a \in A} (X_{\varphi(a)})_{\pi_{a''}('a), \pi_{a'}('a)},$$

$$f_T = \prod_{b \in T_o} (X_{\varphi(b)})_{\pi_{b''}('b), \pi_{b'}('b)} \text{tr}(T_{cl}).$$

We will show that

$$f_0 = \sum_{\tau \in \mathcal{S}_n} \text{sgn}(\tau) f_{T^\tau}. \quad (7)$$

For an open path  $b = b_1 \cdots b_r \in T_o$  with  $b_1, \dots, b_r \stackrel{t}{\in} T$  numbers  $'b_1, \dots, 'b_{r-1}$  are pairwise different. It is not difficult to see that

$$(X_{\varphi(b)})_{\pi_{b''}('b), \pi_{b'}('b)} = \sum_{1 \leq h('b_1), \dots, h('b_{r-1}) \leq n} B(h('b_1), \dots, h('b_{r-1})),$$

where  $B(h('b_1), \dots, h('b_{r-1}))$  stands for

$$(X_{\varphi(b_1)})_{\pi_{b_1''}('b_1), h('b_1)} (X_{\varphi(b_2)})_{h('b_2), \pi_{b_2'}('b_2)} \cdots (X_{\varphi(b_r)})_{h('b_r), \pi_{b_r'}('b_r)}.$$

Similarly, for a closed path  $c = c_1 \cdots c_l \in T_{cl}$  with  $c_1, \dots, c_l \stackrel{cl}{\in} T$  numbers  $'c_1, \dots, 'c_l$  are pairwise different. Therefore

$$\text{tr}(X_{\varphi(c)}) = \sum_{1 \leq h('c_1), \dots, h('c_l) \leq n} C(h('c_1), \dots, h('c_l)),$$

where  $C(h('c_1), \dots, h('c_l))$  stands for

$$(X_{\varphi(c_1)})_{h('c_1), \pi_{c_1'}('c_1)} (X_{\varphi(c_2)})_{\pi_{c_2'}('c_2), h('c_2)} \cdots (X_{\varphi(c_l)})_{\pi_{c_l'}('c_l), h('c_l)}.$$

Summarizing, we obtain

$$f_T = \prod_{b_1 \cdots b_r \in T_o} \sum_{1 \leq h('b_1), \dots, h('b_{r-1}) \leq n} B(h('b_1), \dots, h('b_{r-1}))$$

$$* \prod_{c_1 \cdots c_l \in T_{cl}} \sum_{1 \leq h('c_1), \dots, h('c_l) \leq n} C(h('c_1), \dots, h('c_l)).$$

Since elements of the set  $\{ 'b_i, 'c_j \mid r, l > 0, b_1 \cdots b_r \in T_o, c_1 \cdots c_l \in T_{cl}, 1 \leq i \leq r-1, 1 \leq j \leq l \}$  are pairwise different and their union equals  $[1, n]$ , we infer

$$f_T = \sum_{h: [1, n] \rightarrow [1, n]} \prod_{b_1 \cdots b_r \in T_o, c_1 \cdots c_l \in T_{cl}} B(h('b_1) \cdots h('b_{r-1})) C(h('c_1) \cdots h('c_l)).$$

For an  $a \in T$  and a function  $h : [1, n] \rightarrow [1, n]$  we set:

$$(h, a)^1 = \begin{cases} h('a), & \text{if } a' \in \{q_1, q_2\} \\ \pi_{a'}('a), & \text{otherwise} \end{cases},$$

$$(h, a)^2 = \begin{cases} h(''a), & \text{if } a'' \in \{q_1, q_2\} \\ \pi_{a''}(''a), & \text{otherwise} \end{cases}.$$

Recall that  $\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_{m-2} \in \mathcal{S}_{n_{m-2}}$  have been fixed. Since  $(X_{\varphi(a^t)})_{(h, a^t)^2, (h, a^t)^1} = (X_{\varphi(a)})_{(h, a)^2, (h, a)^1}$  for all  $a \in T$ , we can rewrite  $f_T$  in a form

$$f_T = \sum_{h: [1, n] \rightarrow [1, n]} \prod_{a \in A} (X_{\varphi(a)})_{(h, a)^2, (h, a)^1}.$$

Denote by  $H$  the set of functions  $h : [1, n] \rightarrow [1, n]$  that are not bijections. Since bijections  $h : [1, n] \rightarrow [1, n]$  are in one to one correspondence with permutations from  $\mathcal{S}_n$ , we obtain that

$$\begin{aligned} & \sum_{\tau \in \mathcal{S}_n} \text{sgn}(\tau) f_{T^\tau} \\ &= \sum_{\tau, \pi \in \mathcal{S}_n} \text{sgn}(\tau) \prod_{a \in A} (X_{\varphi(a)})_{(\pi, a^\tau)^2, (\pi, a^\tau)^1} + \sum_{\tau \in \mathcal{S}_n, h \in H} \text{sgn}(\tau) \prod_{a \in A} (X_{\varphi(a)})_{(h, a^\tau)^2, (h, a^\tau)^1}. \end{aligned}$$

Denote the first and the second summands of the last expression by  $f_1$  and  $f_2$ , respectively.

Substituting  $\pi_{m-1}^{-1} \pi_m$  for  $\tau$  and  $\pi_{m-1}$  for  $\pi$  in  $f_1$  we get

$$f_1 = \sum_{\pi_{m-1}, \pi_m \in \mathcal{S}_n} \text{sgn}(\pi_{m-1}) \text{sgn}(\pi_m) \prod_{a \in A} (X_{\varphi(a)})_{(\pi_{m-1}, a^{\pi_{m-1}^{-1} \pi_m})^2, (\pi_{m-1}, a^{\pi_{m-1}^{-1} \pi_m})^1}.$$

The following calculations show that  $(\pi_{m-1}, a^{\pi_{m-1}^{-1} \pi_m})^1 = \pi_{a'}('a)$ .

- If  $a' = q_1 = m-1$ , then  $(\pi_{m-1}, a^{\pi_{m-1}^{-1} \pi_m})^1 = \pi_{m-1}('a^{\pi_{m-1}^{-1} \pi_m}) = \pi_{m-1}('a)$ .

- If  $a' = q_2 = m$ , then  $(\pi_{m-1}, a^{\pi_{m-1}^{-1}\pi_m})^1 = \pi_{m-1}('a^{\pi_{m-1}^{-1}\pi_m}) = \pi_{m-1} \circ \pi_{m-1}^{-1} \circ \pi_m('a) = \pi_m('a)$ .
- If  $a' < m - 1$ , then  $(\pi_{m-1}, a^{\pi_{m-1}^{-1}\pi_m})^1 = \pi_k('a^{\pi_{m-1}^{-1}\pi_m}) = \pi_{a'}('a)$ , where  $k = (a^{\pi_{m-1}^{-1}\pi_m})' = a'$ .

In the same way we obtain  $(\pi_{m-1}, a^{\pi_{m-1}^{-1}\pi_m})^2 = \pi_{a''}('a)$ . Therefore,  $f_1 = f_0$ .

The set  $H$  can be represented as the disjoint union  $H = \bigsqcup_{1 \leq p < q \leq n} H_{pq}$  of sets

$$H_{pq} = \{h \in H \mid h(p) = h(q) \text{ and } h(i) \neq h(j) \text{ for every } (i, j) < (p, q)\},$$

where  $<$  is the lexicographical order on  $\mathbb{N} \times \mathbb{N}$  defined in Section 2. We have

$$f_2 = \sum_{1 \leq p < q \leq n} \sum_{h \in H_{pq}} \left( \sum_{\tau \in \mathcal{S}_n, \tau^{-1}(p) < \tau^{-1}(q)} \text{sgn}(\tau) \prod_{a \in A} (X_{\varphi(a)})_{(h, a^\tau)^2, (h, a^\tau)^1} - \sum_{\tau \in \mathcal{S}_n, \tau^{-1}(p) < \tau^{-1}(q)} \text{sgn}(\tau) \prod_{a \in A} (X_{\varphi(a)})_{(h, a^{(p,q) \circ \tau})^2, (h, a^{(p,q) \circ \tau})^1} \right),$$

where, as usual,  $(p, q)$  stands for a transposition from  $\mathcal{S}_n$ . Compute  $(h, a^{(p,q) \circ \tau})^1$  for  $h \in H_{pq}$ .

- If  $a' = m - 1$ , then  $(h, a^{(p,q) \circ \tau})^1 = h('a) = h('a^\tau)$ .
- If  $a' = m$ , then  $(h, a^{(p,q) \circ \tau})^1 = h \circ (p, q) \circ \tau('a) = h \circ \tau('a) = h('a^\tau)$ .
- If  $a' < m - 1$ , then  $(h, a^{(p,q) \circ \tau})^1 = \pi_{(a^{(p,q) \circ \tau})'}('a^{(p,q) \circ \tau}) = \pi_{a'}('a) = \pi_{a'}('a^\tau)$ .

Hence  $(h, a^{(p,q) \circ \tau})^1 = (h, a^\tau)^1$  and, similarly,  $(h, a^{(p,q) \circ \tau})^2 = (h, a^\tau)^2$ . This implies  $f_2 = 0$  and proves (7).

Rewrite  $\text{bpf}_T^0(X_1, \dots, X_s)$  in a form

$$\sum_{\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_{m-2} \in \mathcal{S}_{n_{m-2}}} \text{sgn}(\pi_1) \cdots \text{sgn}(\pi_{m-2}) \prod_{a \in T, a \notin A} (X_{\varphi(a)})_{\pi_{a''}('a), \pi_{a'}('a)} \cdot f_0.$$

Applying (7), and taking into account that

$$\prod_{b \in T_o^\tau} (X_{\varphi(b)})_{\pi_{b''}('b), \pi_{b'}('b)} = \prod_{a = \tilde{b}, b \in T_o^\tau} (X_{\varphi(a)})_{\pi_{a''}('a), \pi_{a'}('a)},$$

for all  $\tau \in \mathcal{S}_n$  we complete the proof.  $\square$

## 6 Coefficients

Continuing the proof of Theorem 3 we keep the assumptions and notations from previous sections. The aim of this section is to prove formula (14) (see below).

Let  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c})$  be a  $T$ -admissible quadruple with  $\#\underline{b} = p$  and  $\#\underline{c} = q$ . Denote by  $\mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$  the subset of  $\mathcal{S}_n$  containing  $\tau \in \mathcal{S}_n$  if and only if

$$\begin{aligned} \varphi(T_o^\tau) &\stackrel{t}{=} \underbrace{\{b_1, \dots, b_1\}}_{\beta_1}, \dots, \underbrace{\{b_p, \dots, b_p\}}_{\beta_p} \text{ and} \\ \varphi(T_{cl}^\tau) &\stackrel{ct}{=} \{c_1^{\lambda_{11}}, \dots, c_1^{\lambda_{1l_1}}, \dots, c_q^{\lambda_{q1}}, \dots, c_q^{\lambda_{ql_q}}\}_m, \end{aligned}$$

for some positive integers  $\lambda_{i1}, \dots, \lambda_{il_i}$  satisfying  $\lambda_{i1} + \dots + \lambda_{il_i} = \gamma_i$  ( $1 \leq i \leq q$ ), where  $c_i^{\lambda_{ij}}$  stands for  $c_i \cdots c_i$  ( $\lambda_{ij}$  times). Notice that  $\varphi(T_o^\tau)$  and  $\varphi(T_{cl}^\tau)$  are multisets (see Section 4). For  $\tau \in \mathcal{S}_n$  we set

$$\mathcal{S}_\tau = \{\pi \in \mathcal{S}_n \mid \varphi(T_o^\pi) \stackrel{t}{=} \varphi(T_o^\tau), \varphi(T_{cl}^\pi) \stackrel{ct}{=} \varphi(T_{cl}^\tau)\}.$$

The following properties are immediate consequences of the definitions.

**Lemma 5.**

- a) For all  $\pi, \tau \in \mathcal{S}_n$  we have  $\mathcal{S}_\pi = \mathcal{S}_\tau$  or  $\mathcal{S}_\pi \cap \mathcal{S}_\tau = \emptyset$ ;
- b)  $\tau \in \mathcal{S}_\tau$  for all  $\tau \in \mathcal{S}_n$ ;
- c) If  $\tau \in \mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ , then  $\mathcal{S}_\tau \subset \mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ ;
- d)  $\mathcal{S}_n = \bigsqcup_{(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T} \mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ .

Denote by  $\overline{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$  a subset of  $\mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$  such that

$$\mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}} = \bigsqcup_{\tau \in \overline{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}} \mathcal{S}_\tau. \quad (8)$$

The existence of  $\overline{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$  follows from parts a), b), c) of Lemma 5.

Rewrite the coefficient  $c_T = \prod_{j=1}^s \#\{a \in T \mid \varphi(a) = j\}!$  as  $c_T = c_T^{(q_1, q_2)} c_T^{(rest)}$ , where

$$c_T^{(q_1, q_2)} = \prod_{j=1}^s \#\{a \in T \mid \varphi(a) = j, a' > m-2 \text{ or } a'' > m-2\}! \text{ and}$$

$$c_T^{(rest)} = \prod_{j=1}^s \#\{a \in T \mid \varphi(a) = j, a', a'' \leq m-2\}!.$$

Given  $\tau \in \mathcal{S}_n$ , we have  $c_{\tilde{T}\tau} = c_T^{(rest)} c_T^{(\tau)}$ , where

$$c_T^{(\tau)} = \prod_{j=1}^t \#\{b \in T_o^\tau \mid \varphi(\tilde{b}) = j\}!$$

for  $X_{\tilde{T}} = (Y_1, \dots, Y_t)$ . According to part d) of Lemma 5, for every  $\tau \in \mathcal{S}_n$  there is a unique (up to the  $\stackrel{ct}{\sim}$ -equivalence) quadruple  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T$  such that  $\tau \in \mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ . Therefore

$$\varphi(T_{cl}^\tau) \stackrel{ct}{=} \{c_1^{\lambda_{11}}, \dots, c_1^{\lambda_{1l_1}}, \dots, c_q^{\lambda_{q1}}, \dots, c_q^{\lambda_{qlq}}\}_m$$

for some partitions  $\underline{\lambda}_i = (\lambda_{i1}, \dots, \lambda_{il_i}) \vdash \gamma_i$ , where  $1 \leq i \leq q$ . Denote the corresponding multi-partition by

$$\underline{\lambda}(\tau) = (\underline{\lambda}_1, \dots, \underline{\lambda}_q) \vdash \underline{\gamma}. \quad (9)$$

For a vector  $\underline{\lambda} = (\lambda_1, \dots, \lambda_l) \in \mathbb{N}^l$  define  $\underline{\lambda}! = \lambda_1! \cdots \lambda_l!$  and

$$c(\underline{\lambda}) = \lambda_1 \cdots \lambda_l \cdot \prod_{k>0} \#\{1 \leq i \leq l \mid \lambda_i = k\}!. \quad (10)$$

For a multivector  $\underline{\lambda} = (\underline{\lambda}_1, \dots, \underline{\lambda}_q)$  with  $\underline{\lambda}_i \in \mathbb{N}^{l_i}$  we set  $c(\underline{\lambda}) = c(\underline{\lambda}_1) \cdots c(\underline{\lambda}_q)$ .

**Definition (of the  $\sim$ -equivalence).** For  $a_1, \dots, a_p, b_1, \dots, b_q \stackrel{t}{\in} T$  and  $\tau \in \mathcal{S}_n$  consider words  $a = a_1 \cdots a_p$ ,  $b = b_1 \cdots b_q$ , and  $b^\tau = b_1^\tau \cdots b_q^\tau$ . If  $a = b$ , i.e.,  $p = q$  and  $a_1 = b_1, \dots, a_p = b_p$ , then we write  $a \sim b^\tau$  and call  $a$ ,  $b^\tau$  *equivalent*. Analogously, if  $a \stackrel{t}{=} b$  or  $a \stackrel{ct}{=} b$ , respectively, then we write  $a \stackrel{t}{\sim} b^\tau$  or  $a \stackrel{ct}{\sim} b^\tau$ , respectively. We will also use similar notation for multisets.

**Example 5.** Let  $T$  be the tableau

|                 |                 |              |
|-----------------|-----------------|--------------|
| $a$             | $b \rightarrow$ |              |
| $\downarrow$    |                 |              |
| $c$             | $d$             | $e$          |
| $f \rightarrow$ |                 | $\downarrow$ |

of dimension  $(4, 4, 4)$ . Suppose  $q_1 = 1$ ,  $q_2 = 2$ , and  $\tau \in \mathcal{S}_4$  is such a permutation that  $T_o^\tau = \{(d^\tau)^t (a^\tau)^t b^\tau\}_m$  and  $T_{cl}^\tau = \{c^\tau f^\tau\}_m$ . Then  $T_o^\tau \sim \{d^t a^t b\}_m$ ,  $T_{cl}^\tau \sim \{cf\}_m$ , and  $\tau$

is uniquely determined by these data, namely,  $\tau$  is the cycle  $(2, 4, 3)$  (see Lemma 6 and its proof). In particular,  $T^\tau$  is

|              |            |               |
|--------------|------------|---------------|
| $a^\tau$     | $b^\tau$   | $\rightarrow$ |
| $\downarrow$ | $d^\tau$   | $\rightarrow$ |
| $c^\tau$     | $\nearrow$ | $e^\tau$      |
| $f^\tau$     | $\searrow$ | $\downarrow$  |

**Lemma 6.** *Suppose  $\pi_1, \pi_2 \in \mathcal{S}_n$ . Then  $\pi_1 = \pi_2$  if and only if  $T_o^{\pi_1} \stackrel{t}{\sim} T_o^{\pi_2}$  and  $T_{cl}^{\pi_1} \stackrel{ct}{\sim} T_{cl}^{\pi_2}$ .*

*Proof.* **1.** If  $\pi_1 = \pi_2$ , then the statement of lemma is trivial.

**2.** To prove the equality  $\pi_1 = \pi_2$ , it is enough to show that any permutation  $\tau \in \mathcal{S}_n$  is uniquely determined by  $T_o^\tau$ ,  $T_{cl}^\tau$ , and  $T$ .

For any  $1 \leq i \leq n$  there exists an  $a \stackrel{t}{\in} T$ , satisfying  $a' = q_2$  and  $'a = i$ . There is a unique maximal path  $c$  from  $T_o^\tau \sqcup T_{cl}^\tau$  that contains  $a^\tau$  or  $(a^t)^\tau$ . Hence  $c = a_1^\tau \cdots a_k^\tau \cdots a_l^\tau$ , where  $a \stackrel{t}{=} a_k$  for some  $k$ , and  $a_1, \dots, a_l \stackrel{t}{\in} T$ .

Let  $a = a_k$ . If  $k < l$ , then  $(a_{k+1}^\tau)'' = q_1 = a_{k+1}''$  by the definition, which implies  $''a_{k+1}^\tau = ''a_{k+1}$ . Moreover,  $\tau(i) = 'a^\tau = ''a_{k+1}^\tau = ''a_{k+1}$  and  $''a_{k+1}$  is uniquely determined by  $T$ . Therefore in the given case  $\tau(i)$  is uniquely determined by  $T_o^\tau$ ,  $T_{cl}^\tau$ , and  $T$ . If  $k = l$ , then the path  $c$  is open. Taking  $a_1$  instead of  $a_{k+1}$  and acting in the same manner as in the previous case, we obtain that  $\tau(i) = ''a_1$ .

The case of  $a = a_k^t$  can be treated analogously.  $\square$

**Lemma 7.** *Let  $a, b \stackrel{t}{\in} T$  and  $a \neq b$ .*

a) *Suppose  $a' = b' \in \{q_1, q_2\}$ ,  $a'' = b'' \in \{q_1, q_2\}$ . Then for  $\pi = ('a, ''b) \circ ('a, 'b) \in \mathcal{S}_n$  we have*

$$T_o^\pi \stackrel{t}{\sim} T_o|_{a \rightarrow b, b \rightarrow a}, \quad T_{cl}^\pi \stackrel{ct}{\sim} T_{cl}|_{a \rightarrow b, b \rightarrow a}.$$

b) *Suppose  $a' = b' \in \{q_1, q_2\}$ ,  $a'' = b'' \notin \{q_1, q_2\}$ . Then for the transposition  $\pi = ('a, 'b) \in \mathcal{S}_n$  we have*

$$T_o^\pi \stackrel{t}{\sim} T_o|_{a \rightarrow b, b \rightarrow a}, \quad T_{cl}^\pi \stackrel{ct}{\sim} T_{cl}.$$

*Proof.* **a)** Let  $a' = b' = q_1$ ,  $a'' = b'' = q_2$ . Moreover, assume that  $'a, ''a, 'b, ''b$  are pairwise different. Then  $a, b$  belong to some maximal paths  $a_0, b_0$  in  $T$ , i.e.,  $a_0 = a_1 \cdots a_2 a a_3 \cdots a_4$  and  $b_0 = b_1 \cdots b_2 b b_3 \cdots b_4$ , where  $a_i, b_j \stackrel{t}{\in} T$ . It is not difficult to see that maximal paths in  $T^\pi$ , containing  $a^\pi$  or  $b^\pi$ , are  $a_1^\pi \cdots a_2^\pi b^\pi a_3^\pi \cdots a_4^\pi$  and  $b_1^\pi \cdots b_2^\pi a^\pi b_3^\pi \cdots b_4^\pi$ . The remaining maximal paths in  $T^\pi$  are equivalent to paths in  $T$ . The claim follows.

Other possibilities for  $a, b$  can be treated in the similar fashion.

**b)** Analogous to part a).  $\square$

Recall that Lemma 4 states that  $\text{bpf}_T^0(X_1, \dots, X_s) = \sum_{\tau \in \mathcal{S}_n} g(T, \tau)$ , where

$$g(T, \tau) = \text{sgn}(\tau) \text{bpf}_{\tilde{T}^\tau}^0(X_{\tilde{T}^\tau}) \text{tr}(T_{cl}^\tau). \quad (11)$$

**Lemma 8.** *Let  $\tau \in \mathcal{S}_n$ ,  $a, b \stackrel{t}{\in} T^\tau$ ,  $a \neq b$ , and  $\varphi(a) = \varphi(b)$ . If  $a' = b' \in \{q_1, q_2\}$  and  $a'' = b'' \notin \{q_1, q_2\}$ , then  $g(T, ('a, 'b) \circ \tau) = g(T, \tau)$ .*

*Proof.* **a)** Let  $a' = b' = q_1$ . For simplicity assume  $\tau = id$  is the identical permutation. Consider paths  $u = u_1 \cdots u_l, v$  in  $T$  such that  $au, bv \stackrel{t}{\in} T_o$ ,  $u_1, \dots, u_l \stackrel{t}{\in} T$  and set  $k = a'' = b''$  and  $\xi = ('a, 'b) \in \mathcal{S}_n$ . There are two cases.

**Case**  $u_l \neq b^t$ . We have

$$T_o^\xi \stackrel{t}{\sim} T_o \bigcup \{a^\xi v^\xi, b^\xi u^\xi\} \stackrel{t}{\setminus} \{au, bv\}, \quad T_{cl}^\xi \stackrel{ct}{\sim} T_{cl}.$$

Therefore

$$g(T, \xi) = - \sum_{\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_{m-2} \in \mathcal{S}_{n_{m-2}}} \text{sgn}(\pi_1) \cdots \text{sgn}(\pi_{m-2}) \text{tr}(T_{cl}) * \\ \prod_{d \in \tilde{T}, d \neq \widetilde{au}, \widetilde{bv}} (X_{\varphi(d)})_{\pi_{d''}(''d), \pi_{d'}('d)} \cdot (X_{\varphi(av)})_{\pi_k(''a), \pi_{v'}('v)} (X_{\varphi(bu)})_{\pi_k(''b), \pi_{u'}('u)}.$$

Observe that  $X_{\varphi(av)} = X_{\varphi(bv)}$  and  $X_{\varphi(bu)} = X_{\varphi(au)}$  and substitute  $\pi_k \circ (''a, ''b)$  for  $\pi_k$ . Then  $\pi_k(''a)$  turns into  $\pi_k(''b)$ ,  $\pi_k(''b)$  turns into  $\pi_k(''a)$ , and the rest of  $\pi_p(q)$  does not change for  $1 \leq p \leq m-2$ ,  $q \in \mathcal{S}_{n_p}$ . This proves the claim.

**Case**  $u_l = b^t$ . Denote  $u_1 \cdots u_{l-1}$  by  $w$ . Then  $awb^t \stackrel{t}{\in} T_o$  and

$$T_o^\xi \stackrel{t}{\sim} T_o \bigcup \{a^\xi (w^\xi)^t (b^\xi)^t\} \stackrel{t}{\setminus} \{awb^t\}, \quad T_{cl}^\xi \stackrel{ct}{\sim} T_{cl}.$$

Therefore

$$g(T, \xi) = - \sum_{\pi_1 \in \mathcal{S}_{n_1}, \dots, \pi_{m-2} \in \mathcal{S}_{n_{m-2}}} \text{sgn}(\pi_1) \cdots \text{sgn}(\pi_{m-2}) \text{tr}(T_{cl})^* \\ \prod_{d \in \widetilde{T}, d \neq awb^t} (X_{\varphi(d)})_{\pi_d''(''d), \pi_d'(''d)} \cdot (X_{\varphi(aw^t b^t)})_{\pi_k(''a), \pi_k(''b)}.$$

Observe that  $X_{\varphi(aw^t b^t)} = X_{\varphi(awb^t)}^t$ . Substitution of  $\pi_k \circ (''a, ''b)$  for  $\pi_k$  completes the proof.

The proof for an arbitrary  $\tau$  is exactly the same as for  $\tau = id$ .

b) The proof in the case  $a' = b' = q_2$  is analogous.  $\square$

**Lemma 9.** *Suppose  $\tau \in \mathcal{S}_n$ . Then*

$$a) \# \mathcal{S}_\tau = \frac{c_T^{(q_1, q_2)}}{\beta! c(\underline{\lambda}(\tau))}, \text{ where } \tau \in \mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}, (\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in \mathcal{Q}_T.$$

$$b) \text{ If } \pi \in \mathcal{S}_\tau, \text{ then } g(T, \pi) = g(T, \tau).$$

*Proof.* a) Given  $\pi_1, \pi_2 \in \mathcal{S}_n$ , we write  $(T_o^{\pi_1}, T_{cl}^{\pi_1}) \stackrel{ct}{\sim} (T_o^{\pi_2}, T_{cl}^{\pi_2})$  and say that these pairs are equivalent if  $T_o^{\pi_1} \stackrel{t}{\sim} T_o^{\pi_2}$  and  $T_{cl}^{\pi_1} \stackrel{ct}{\sim} T_{cl}^{\pi_2}$ . As in Section 5 define

$$A = \{a \in T^\tau \mid a' > m - 2 \text{ or } a'' > m - 2\}.$$

Let  $S$  be the set consisting of permutations  $\pi \in \mathcal{S}_n$  for which there is a bijection  $\Lambda : A \rightarrow A$  satisfying

$$\varphi(\Lambda(a)) = \varphi(a) \text{ for all } a \in A, \quad (12)$$

$$(\overline{\Lambda}(T_o^\tau), \overline{\Lambda}(T_{cl}^\tau)) \stackrel{ct}{\sim} (T_o^\pi, T_{cl}^\pi), \quad (13)$$

where  $\overline{\Lambda}(a) = \Lambda(a)$ ,  $\overline{\Lambda}(a^t) = \Lambda(a)^t$  for an  $a \in A$ , and  $\overline{\Lambda}(a_1 \cdots a_l) = \overline{\Lambda}(a_1) \cdots \overline{\Lambda}(a_l)$  for  $a_1, \dots, a_l \stackrel{t}{\in} A$ .

Definitions of  $S$  and  $\mathcal{S}_\tau$  imply the inclusion  $S \subset \mathcal{S}_\tau$ . On the other hand, it is not difficult to see that  $\mathcal{S}_\tau \subset S$ . Thus  $\mathcal{S}_\tau = S$ .

A bijection  $\Lambda : A \rightarrow A$  can be considered as a permutation of  $A$  and can be written as a composition of transpositions on  $A$ . By Lemma 7, for every bijection  $\Lambda : A \rightarrow A$  satisfying (12) there exists a  $\pi \in \mathcal{S}_n$  such that (13) holds. This and Lemma 6 show that  $\#S$  equals the number of  $\stackrel{ct}{\sim}$ -equivalence classes of pairs

$(\bar{\Lambda}(T_o^\tau), \bar{\Lambda}(T_{cl}^\tau))$  for bijections  $\Lambda : A \rightarrow A$  satisfying (12). Combinatorial argument completes the proof.

**b)** Using the notation of part a), assume  $\pi \in \mathcal{S}_\tau$ . Since  $\mathcal{S}_\tau = S$ , there exists a bijection  $\Lambda : A \rightarrow A$  satisfying (12) and (13). As in part a), for such  $\Lambda$  we construct  $\pi_0 \in \mathcal{S}_n$  such that

$$(\bar{\Lambda}(T_o^\tau), \bar{\Lambda}(T_{cl}^\tau)) \stackrel{ct}{\sim} (T_o^{\pi_0 \circ \tau}, T_{cl}^{\pi_0 \circ \tau})$$

and  $\pi_0$  is a composition of permutations of types

1.  $(''a, ''b) \circ ('a, 'b)$ ,
2.  $('a, 'b)$ ,

where  $a, b \stackrel{t}{\in} T^\tau$ ,  $\varphi(a) = \varphi(b)$  (apply parts a), b) of Lemma 7 to  $T^\tau$ ). Using Lemma 6 we infer  $\pi = \pi_0 \circ \tau$ . For permutations of the first type we obtain  $g(T, (''a, ''b) \circ ('a, 'b) \circ \tau) = g(T, \tau)$ ; for permutations of the second type we use Lemma 8.  $\square$

Now we can express  $\text{bpf}_T(X_1, \dots, X_s)$  in a suitable form. By Lemma 4,  $\text{bpf}_T(X_1, \dots, X_s) = \frac{1}{c_T} \sum_{\tau \in \mathcal{S}_n} g(T, \tau)$ . Using Lemma 5 and notation from the beginning of this section, we obtain

$$\text{bpf}_T(X_1, \dots, X_s) = \frac{1}{c_T} \sum_{(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T} \sum_{\pi \in \bar{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}} \sum_{\tau \in \mathcal{S}_\pi} g(T, \tau).$$

Lemma 9 yields

$$\begin{aligned} & \text{bpf}_T(X_1, \dots, X_s) \\ &= \frac{1}{c_T^{(q_1, q_2)} c_T^{(rest)}} \sum_{(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T} \sum_{\pi \in \bar{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}} \frac{c_T^{(q_1, q_2)}}{\underline{\beta}! c(\underline{\lambda}(\pi))} \text{sgn}(\pi) \text{bpf}_{\tilde{T}^\pi}^0(X_{\tilde{T}^\pi}) \text{tr}(T_{cl}^\pi). \end{aligned}$$

Let us recall that  $c_{\tilde{T}^\pi} = c_T^{(rest)} c_T^{(\pi)}$ . Since  $c_T^{(\pi)} = \underline{\beta}!$  for  $\pi \in \bar{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$  we conclude

$$\text{bpf}_T(X_1, \dots, X_s) = \sum_{(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T} \sum_{\pi \in \bar{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}} \text{sgn}(\pi) \frac{1}{c(\underline{\lambda}(\pi))} \text{tr}(T_{cl}^\pi) \text{bpf}_{\tilde{T}^\pi}(X_{\tilde{T}^\pi}). \quad (14)$$

## 7 Traces and $\sigma_k$

Continue working under the same assumptions as in the previous section.

For  $\gamma \in \mathbb{N}$ ,  $\underline{\nu} = (\nu_1, \dots, \nu_l) \vdash \gamma$  denote by  $\rho_{\underline{\nu}} \in \mathcal{S}_\gamma$  the composition of cycles

$$\rho_{\underline{\nu}} = (1, 2, \dots, \nu_1) \circ (\nu_1 + 1, \dots, \nu_1 + \nu_2) \circ \dots \circ (\nu_1 + \dots + \nu_{l-1}, \dots, \gamma).$$

Note that  $\text{sgn}(\rho_{\underline{\nu}}) = (-1)^{\gamma - \#\underline{\nu}}$ .

Let  $(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T$ ,  $\xi = \xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ , and  $\#\underline{\gamma} = q$ . Without loss of generality we can assume that

$$\begin{aligned} T_{cl}^\xi &= \{d_{11}, \dots, d_{1\gamma_1}, \dots, d_{q1}, \dots, d_{q\gamma_q}\}_m, \quad \text{where} \\ d_{ij} &\in \mathcal{M}_T, \quad \varphi(d_{ij}) = c_i, \quad d''_{ij} = q_1, \quad d'_{ij} = q_2, \quad {}''d_{ij} = {}'d_{ij} \end{aligned} \quad (15)$$

for any  $1 \leq i \leq q$  and  $1 \leq j \leq \gamma_i$ . There is a unique bijection  $\phi_i : [1, \gamma_i] \rightarrow \{{}'d_{i1}, \dots, {}'d_{i\gamma_i}\}$  that is a monotone increasing map, i.e.,  $\phi_i(j_1) < \phi_i(j_2)$  for  $j_1 < j_2$ .

Let  $S = \{\pi_{1, \underline{\nu}_1} \circ \dots \circ \pi_{q, \underline{\nu}_q} \circ \xi \mid \underline{\nu}_1 \vdash \gamma_1, \dots, \underline{\nu}_q \vdash \gamma_q\}$ , where the permutation  $\pi_{i, \underline{\nu}_i} \in \mathcal{S}_n$  is defined by

$$\pi_{i, \underline{\nu}_i}(k) = \begin{cases} \phi_i \circ \rho_{\underline{\nu}_i} \circ \phi_i^{-1}(k), & \text{if } k \in \text{Im } \phi_i \\ k, & \text{otherwise} \end{cases}.$$

Then

- By (9),  $\underline{\lambda}(\pi_{1, \underline{\nu}_1} \circ \dots \circ \pi_{q, \underline{\nu}_q} \circ \xi) = (\underline{\nu}_1, \dots, \underline{\nu}_q)$  is a multi-partition of  $\underline{\gamma}$ .
- Suppose  $\pi, \tau \in \mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}$ . Then  $\mathcal{S}_\tau = \mathcal{S}_\pi$  if and only if  $\underline{\lambda}(\tau) = \underline{\lambda}(\pi)$ .

These remarks imply that  $\mathcal{S}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}} = \bigsqcup_{\tau \in S} \mathcal{S}_\tau$ , hence without loss of generality we can assume that  $\overline{\mathcal{S}}_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}} = S$  (see (8)).

Note that

- Since  $\text{sgn}(\pi_{i, \underline{\nu}_i}) = \text{sgn}(\rho_{\underline{\nu}_i})$ , we have the equality  $\text{sgn}(\pi_{1, \underline{\nu}_1} \circ \dots \circ \pi_{q, \underline{\nu}_q} \circ \xi) = \text{sgn}(\rho_{\underline{\nu}_1}) \cdots \text{sgn}(\rho_{\underline{\nu}_q}) \text{sgn}(\xi)$ .
- $\text{tr}(T_{cl}^{\pi_{1, \underline{\nu}_1} \circ \dots \circ \pi_{q, \underline{\nu}_q} \circ \xi}) = \prod_{i=1}^q \prod_{j=1}^{\#\underline{\nu}_i} \text{tr}(X_{c_i}^{\nu_{ij}})$ .
- Since  $T_o^\eta \stackrel{t}{=} T_o^\xi$  for  $\eta = \pi_{1, \underline{\nu}_1} \circ \dots \circ \pi_{q, \underline{\nu}_q} \circ \xi$ , it is easy to see that  $(\widetilde{T}^\eta, X_{\widetilde{T}^\eta}) = (\widetilde{T}^\xi, X_{\widetilde{T}^\xi})$ .

Therefore equality (14) yields  $\text{bpf}_T(X_1, \dots, X_s) =$

$$\sum_{(\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}) \in Q_T} \text{sgn}(\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}) \text{bpf}_{\tilde{T}^{\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}}}(X_{\tilde{T}^{\xi_{\underline{\beta}, \underline{\gamma}, \underline{b}, \underline{c}}}}) \prod_{i=1}^{\#\underline{\gamma}} \left( \sum_{\underline{\nu} \vdash \gamma_i} (-1)^{\gamma_i - \#\underline{\nu}} \frac{1}{c(\underline{\nu})} \prod_{j=1}^{\#\underline{\nu}} \text{tr}(X_{c_i}^{\nu_j}) \right). \quad (16)$$

This formula together with the following well known lemma completes the proof of Theorem 3.

**Lemma 10.** *Assume  $K = \mathbb{Q}$ . If  $k, n \in \mathbb{N}$ ,  $0 < k \leq n$ , and  $c$  is an  $n \times n$  matrix, then*

$$\sigma_k(c) = \sum_{\underline{\nu} \vdash k} (-1)^{k - \#\underline{\nu}} \frac{1}{c(\underline{\nu})} \prod_{j=1}^{\#\underline{\nu}} \text{tr}(c^{\nu_j}),$$

where  $c(\underline{\nu})$  is given by (10).

Applying (16) for the tableau  $T$  from part 3 of Example 2 we obtain an alternative proof of this lemma.

## 8 Corollaries

The decomposition formula generalizes Amitsur's formula for the determinant (see [1]).

**Corollary 11. (Amitsur).** *For  $k_1 + \dots + k_s = n$  and  $n \times n$  matrices  $X_1, \dots, X_s$  we have*

$$\det_{k_1, \dots, k_s}(X_1, \dots, X_s) = \sum (-1)^{n + (i_1 + \dots + i_q)} \sigma_{i_1}(c_1) \dots \sigma_{i_q}(c_q),$$

where the sum ranges over all  $\stackrel{c}{=}$ -equivalent classes of multisets

$\{\underbrace{c_1, \dots, c_1}_{i_1}, \dots, \underbrace{c_q, \dots, c_q}_{i_q}\}_m$  such that

- $c_1, \dots, c_q$  are primitive words in letters  $X_1, \dots, X_s$  such that  $c_1, \dots, c_q$  are pairwise different with respect to  $\stackrel{c}{=}$ ;
- $i_1 \deg_{X_j} c_1 + \dots + i_q \deg_{X_j} c_q = n_j$  for any  $1 \leq j \leq s$ .

*Proof.* Consider the tableau with substitution  $(T, (X_1, \dots, X_s))$  of dimension  $(n, n)$  from part 2 of Example 2 and assume  $\#\{a \in T \mid \varphi(a) = j\} = k_j$  for any  $1 \leq j \leq s$ . Then  $\text{bpf}_T(X_1, \dots, X_s) = \det_{k_1, \dots, k_s}(X_1, \dots, X_s)$  and the claim follows easily from the decomposition formula.  $\square$

**Corollary 12.** *Let  $Y, Z$  be  $n \times n$  matrices and  $n$  is even. Then  $\overline{\text{pf}}(Y)\overline{\text{pf}}(Z) =$*

$$\sum (-1)^{i_1(\deg_Y c_1 + \deg_Z c_1 + 1) + \dots + i_q(\deg_Y c_q + \deg_Z c_q + 1)} \sigma_{i_1}(c_1) \cdots \sigma_{i_q}(c_q),$$

where the sum ranges over  $\stackrel{ct}{=}$ -equivalent classes of multisets  $\{\underbrace{c_1, \dots, c_1}_{i_1}, \dots, \underbrace{c_q, \dots, c_q}_{i_q}\}_m$  such that

- $c_1, \dots, c_q$  are primitive words in letters  $Y, Y^t, Z, Z^t$  such that  $c_1, \dots, c_q$  are pairwise different with respect to  $\stackrel{ct}{=}$ ;
- words  $c_1, \dots, c_q$  are products of the words  $AB$ , where  $A \in \{Y, Y^t\}$  and  $B \in \{Z, Z^t\}$ .
- $i_1(\deg_Y c_1 + \deg_{Y^t} c_1) + \dots + i_q(\deg_Y c_q + \deg_{Y^t} c_q) = n/2$ ,  
 $i_1(\deg_Z c_1 + \deg_{Z^t} c_1) + \dots + i_q(\deg_Z c_q + \deg_{Z^t} c_q) = n/2$ .

*Proof.* Consider the tableau with substitution  $(T, (Y, Z))$  of dimension  $(n, n)$  from part 4 of Example 2, where  $t = 0, r = s = n/2$ . Then  $\text{bpf}_T(Y, Z) = \overline{\text{pf}}(Y)\overline{\text{pf}}(Z)$  and the formula follows from Theorem 3.  $\square$

**Example 6.** 1. Let  $Y, Z$  be  $2 \times 2$  matrices. Then

$$\overline{\text{pf}}(Y)\overline{\text{pf}}(Z) = \text{tr}(YZ^t) - \text{tr}(YZ).$$

2. Suppose  $Y, Z$  are  $4 \times 4$  matrices. Then  $\overline{\text{pf}}(Y)\overline{\text{pf}}(Z) =$

$$\sigma_2(YZ) + \sigma_2(YZ^t) - \text{tr}(YZ)\text{tr}(YZ^t) + \text{tr}(YZYZ^t) + \text{tr}(YZY^tZ) - \text{tr}(YZY^tZ^t).$$

**Remark 13.** *In the same manner as in Corollary 12 we can also write a formula for a product of two partial linearizations of pfaffians.*

**Acknowledgements.** The first version of this paper was written during author's visit to University of Antwerp, sponsored by Marie Curie Research Training Network Liegrits. The author is grateful for this support. The author would like to thank Fred Van Oystaeyen for his hospitality. This research was also supported by RFFI 05-01-00057.

## References

- [1] S.A. Amitsur, *On the characteristic polynomial of a sum of matrices*, Linear and Multilinear Algebra **8** (1980), 177–182.
- [2] S. Donkin, *Invariants of several matrices*, Invent. Math. **110** (1992), 389–401.
- [3] A.A. Lopatin, A.N. Zubkov, *Semi-invariants of mixed representations of quivers*, Transform. Groups **12** (2007), No. 2, 341–369.
- [4] A.A. Lopatin, *Invariants of quivers under the action of classical groups*, arXiv: math.RT/0608750.
- [5] A.A. Lopatin, *Relations between  $O(n)$ -invariants of several matrices*, in preparation.
- [6] C. Procesi, *The invariant theory of  $n \times n$  matrices*, Adv. Math. **19** (1976), 306–381.
- [7] A.N. Zubkov, *A generalization of the Razmyslov–Procesi theorem*, Algebra and Logic **35** (1996), No. 4, 241–254.
- [8] A.N. Zubkov, *Invariants of an adjoint action of classical groups*, Algebra and Logic **38** (1999), No. 5, 299–318.
- [9] A.N. Zubkov, *Invariants of mixed representations of quivers I*, J. Algebra Appl., **4** (2005), No. 3, 245–285.
- [10] A.N. Zubkov, *Invariants of mixed representations of quivers II: Defining relations and applications*, J. Algebra Appl., **4** (2005), No. 3, 287–312.